

2006 美國病人安全年度目標

(2006 Joint Commission National Patient Safety Goals)

Goal 1 Improve the accuracy of patient identification.

提升病人辨識的正確性

Goal 2 Improve the effectiveness of communication among caregivers.

提升醫護人員間溝通的有效性

Goal 3 Improve the safety of using medications.

改善用藥的安全性

Goal 4 Eliminate wrong-site, wrong-patient, wrong-procedure surgery.

消除手術部位錯誤、病人錯誤、術式錯誤

Goal 5 Retired in 2006. [Improve the safety of using infusion pumps.]

2006 年已取消（改善輸液幫浦的使用安全）

Goal 6 Improve the effectiveness of clinical alarm systems.

提升臨床警示系統的有效性

Goal 7 Reduce the risk of health care-associated infections.

減少健康照護相關的感染風險

Goal 8 Accurately and completely reconcile medications across the continuum

確保病人持續性照護之用藥正確性及完整性

Goal 9 Reduce the risk of patient harm resulting from falls.

減少病人因跌倒造成傷害的風險

Goal 10 Reduce the risk of influenza and pneumococcal disease in older adults.

降低老年人罹患流行性感冒及肺炎的風險

Goal 11 Reduce the risk of surgical fires.

降低手術失火的風險

Goal 12 Implement applicable National Patient Safety Goals and Requirements at the component and practitioner site levels.

基層醫療院所層級需要執行適當的病人安全目標與要求

Goal 13 Encourage the active involvement of residents and their families in the resident's care as a resident safety strategy.

病人安全策略應含鼓勵病人及其家屬主動參與病患照護

Goal 14 Prevent health care-associated pressure ulcers (decubitus ulcers).

避免因健康照護造成的壓瘡